

# IMOXI™

Topical Solution  
(imidacloprid + moxidectin)



**COVER THEM WITH DESIGNER FLEA,  
HEARTWORM AND INTESTINAL  
PARASITE PROTECTION WITHOUT  
PAYING DESIGNER PRICES.**



Fleas



Heartworms



Intestinal  
Parasites



Ear Mites  
(cats only)

**vetoquinol**  
ACHIEVE MORE TOGETHER

**IMOXI™ (IMIDACLOPRID +  
MOXIDECTIN) TOPICAL SOLUTION  
PROVIDES THE SAME MONTHLY  
HEARTWORM, FLEA AND  
INTESTINAL PARASITE PROTECTION  
AS ADVANTAGE MULTI®  
(IMIDACLOPRID + MOXIDECTIN), BUT  
AT A MORE ATTRACTIVE PRICE.**

IMOXI™ has the same antiparasitic formulation used at animal shelters nationwide to treat dogs and cats.

And IMOXI™ comes from Vetoquinol, a trusted family-owned manufacturer dedicated exclusively to animal health since 1933, backing its products with a 100% Satisfaction Guarantee. If you are not satisfied, contact your IMOXI retailer or call 1-817-529-7500 or visit [www.vetoquinolusa.com/satisfaction-guarantee](http://www.vetoquinolusa.com/satisfaction-guarantee).

**APPLY MONTHLY. PROTECT YEARLY.**

Monthly applications are important, because fleas, heartworms and intestinal parasites are year-round threats. One easy-to-apply application of IMOXI™ each month gives your pet year-round protection, and you, year-round peace of mind.

# DOGS AND CATS ENCOUNTER ALL SORTS OF PARASITES, INDOORS AND OUTDOORS.



## Fleas

Fleas are found outdoors and inside and cause your pets to scratch, chew, bite and become restless. Fleas you can see are only the tip of the iceberg. Their family can infest your home, living in carpets, bedding, furniture and flooring. When looking for their next blood meal, they will bite you and your pet!



## Heartworms

While most common in dogs, heartworms can be found in cats. They are transmitted through infected mosquitoes and cause coughing, sluggishness and difficult breathing. Heartworms can be fatal; there is no treatment for heartworms in cats.



## Intestinal Parasites

Intestinal parasites cause inflammation, anemia, diarrhea and even death in some puppies and kittens.

**Some intestinal parasites can be a danger to your family! These zoonotic diseases can be transmitted from animals to humans. Anyone coming in contact with fecal-contaminated soil is at risk, especially infants, the elderly or those with compromised immune systems.**



## Ear Mites – only for cats

Ear mites can be picked up from the environment or other pets. Signs include shaking of the head, ear scratching, inflammation of the ear canals and large accumulations of earwax.

**PROTECT  
YOUR FURRY  
FASHIONISTAS  
WITH THE SAME  
HEARTWORM,  
FLEA AND  
INTESTINAL  
PARASITE  
PROTECTION  
TRUSTED BY  
SHELTERS  
NATIONWIDE.**



**Approved for use in dogs and puppies at least 7 weeks of age and weighing at least 3 lbs.**

**IMOXI™** Topical Solution for Dogs  
(imidacloprid + moxidectin) **is indicated for:**

- Prevention of heartworm disease
- Killing adult fleas on contact—no bite necessary
- Treatment of flea infestations
- Treatment and control of sarcoptic mange
- Treatment and control of intestinal parasites, including hookworm, roundworm and whipworm



**Approved for use in cats and kittens 9 weeks of age and older and weighing at least 2 lbs.**

**IMOXI™** Topical Solution for Cats  
(imidacloprid + moxidectin) **is indicated for:**

- Prevention of heartworm disease
- Killing adult fleas on contact—no bite necessary
- Treatment of flea infestations
- Treatment and control of ear mite infestations
- Treatment and control of intestinal parasites, including hookworm and roundworm

# IMOXI™ (imidacloprid + moxidectin)

Topical Solution is easy to apply. Start by choosing the correct product and dose for your dog or cat.

Dog (lbs.)	Volume (mL)
3-9	0.4
9.1-20	1.0
20.1-55	2.5
55.1-88	4.0
88.1-110*	5.0

\*Dogs over 110 lbs. should be treated with the appropriate combination of IMOXI™ (imidacloprid + moxidectin) Topical Solution for Dogs tubes.

Cat (lbs.)	Volume (mL)
2-5	0.23
5.1-9	0.4
9.1-18*	0.8

\*Cats over 18 lbs. should be treated with the appropriate combination of IMOXI™ (imidacloprid + moxidectin) Topical Solution for Cats tubes.



eliminates a step in the application process.



1. Remove one dose applicator tube from the package. Administer the entire contents of the IMOXI™ Topical Solution for Dogs and Cats tube that correctly corresponds with the body weight of the pet. See Product Insert for complete application instructions.
2. Hold the Twist-N-Go™ tube in an upright position and twist the tip clockwise about ½ turn to break the tube's seal. Remove the cap from the tube.
3. **For Cats:** Part the hair on the back of the cat's neck at the base of the head, until the skin is visible.  
**For Dogs:** The dog should be standing for application. Position the dispensing tip on the dog's back between the shoulder blades. The dispensing tip of the tube can be used to part the dog's hair until the skin is visible.
4. Place the tip of the tube on the skin and apply the entire contents directly on the exposed skin. Lift the tube away from the skin before releasing pressure on the tube.

Do not get this product in your pet's mouth or eyes or allow them to lick the application site for 30 minutes. Treatment at the base of the head will minimize the opportunity for ingestion by grooming. In households with multiple pets, keep animals separated to prevent licking of the application site.

## CAUTION

Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian. Dogs: **WARNING: DO NOT ADMINISTER THIS PRODUCT ORALLY.** For the first 30 minutes after application, ensure that dogs cannot lick the product from application sites on themselves or other treated animals. Children should not come in contact with the application sites for two (2) hours after application. (See Contraindications, Warnings, Human Warnings and Adverse Reactions for more information.) Cats: Do not use on sick, debilitated, or underweight cats. Avoid oral ingestion. For full prescribing information, see back of document.

# IMOXI™ Topical Solution for Dogs and for Cats (imidacloprid + moxidectin)

## BRIEF SUMMARY

Before using IMOXI™ Topical Solution for Dogs (imidacloprid + moxidectin) or IMOXI™ Topical Solution for Cats (imidacloprid + moxidectin), please consult the product insert, a summary of which follows:

## CAUTION

Federal (U.S.A.) Law restricts this drug to use by or on the order of a licensed veterinarian.

## WARNING

### DO NOT ADMINISTER THIS PRODUCT ORALLY

For the first 30 minutes after application ensure that dogs cannot lick the product from application sites on themselves or other treated animals. Children should not come in contact with application sites for two (2) hours after application. (See Contraindications, Warnings, Human Warnings, and Adverse Reactions, for more information.)

## INDICATIONS

**IMOXI™ Topical Solution for Dogs** is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis* and the treatment of *Dirofilaria immitis* circulating microfilariae in heartworm-positive dogs. IMOXI™ Topical Solution for Dogs kills adult fleas and is indicated for the treatment of flea infestations (*Ctenocephalides felis*). IMOXI™ Topical Solution for Dogs is indicated for the treatment and control of sarcoptic mange caused by *Sarcoptes scabiei* var. *canis*. IMOXI™ Topical Solution for Dogs is also indicated for the treatment and control of the following intestinal parasites species: Hookworms (*Ancylostoma caninum*) (*Uncinaria stenocephala*), Roundworms (*Toxocara canis*) (*Toxascaris leonina*) and Whipworms (*Trichuris vulpis*).

**IMOXI™ Topical Solution for Cats** is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis*. IMOXI™ Topical Solution for Cats kills adult fleas (*Ctenocephalides felis*) and is indicated for the treatment of flea infestations. IMOXI™ Topical Solution for Cats is also indicated for the treatment and control of ear mite (*Otodectes cynotis*) infestations and the intestinal parasites species Hookworm (*Ancylostoma tubaeforme*) and Roundworm (*Toxocara cati*).

## CONTRAINDICATIONS

Do not administer this product orally. (See WARNINGS)  
Do not use the Dog product (containing 2.5% moxidectin) on cats.

## WARNINGS

**IMOXI™ Topical Solution for Dogs:** For the first 30 minutes after application: Ensure that dogs cannot lick the product from application sites on themselves or other treated dogs, and separate treated dogs from one another and from other pets to reduce the risk of accidental ingestion. Ingestion of this product by dogs may cause serious adverse reactions including depression, salivation, dilated pupils, incoordination, panting, and generalized muscle tremors. In avermectin sensitive dogs,<sup>a</sup> the signs may be more severe and may include coma and death.<sup>b</sup>

<sup>a</sup> Some dogs are more sensitive to avermectins due to a mutation in the MDR1 gene. Dogs with this mutation may develop signs of severe avermectin toxicity if they ingest this product. The most common breeds associated with this mutation include Collies and Collie crosses.

<sup>b</sup> Although there is no specific antagonist for avermectin toxicity, even severely affected dogs have completely recovered from avermectin toxicity with intensive veterinary supportive care.

**IMOXI™ Topical Solution for Cats:** Do not use on cats less than 9 weeks of age or less than 2 lbs. body weight.

## HUMAN WARNINGS

Not for human use. Keep out of the reach of children. Dogs: Children should not come in contact with the application sites for two (2) hours after application. Cats: Children should not come into contact with the application site for 30 minutes after application. Causes eye irritation. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Exposure to the product has been reported to cause headache; dizziness; and redness, burning, tingling, or numbness of the skin.

### Wash hands thoroughly with soap and warm water after handling.

If contact with eyes occurs, hold eyelids open and flush with copious amounts of water for 15 minutes. If eye irritation develops or persists, contact a physician. If swallowed, call poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. People with known hypersensitivity to benzyl alcohol, imidacloprid or moxidectin should administer the product with caution. In case of allergic reaction, contact a physician. If contact with skin or clothing occurs, take off contaminated clothing. Wash skin immediately with plenty of soap and water. Call a poison control center or physician for treatment advice. The Safety Data Sheet (SDS) provides additional occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Vetoquinol USA at 1-800-835-9496.

## PRECAUTIONS

Do not dispense dose applicator tubes without complete safety and administration information. Use with caution in sick, debilitated, or underweight animals. The safety of IMOXI™ Topical Solution for Dogs has not been established in breeding, pregnant, or lactating dogs. The safe use of IMOXI™ Topical Solution for Dogs has not been established in puppies and dogs less than 7 weeks of age or less than 3 lbs. body weight. Cats may experience hypersalivation, tremors, vomiting and decreased appetite if IMOXI™ Topical Solution for Cats is inadvertently administered orally or through grooming/licking of the application site. The safety of IMOXI™ Topical Solution for Cats has not been established in breeding, pregnant, or lactating cats. The effectiveness of IMOXI™ Topical Solution for Cats against heartworm infections (*D. immitis*) after bathing has not been evaluated in cats. Use of this product in geriatric patients with subclinical conditions has not been adequately studied. Several otherwise healthy, thin geriatric cats experienced prolonged lethargy and sleepiness after using imidacloprid and moxidectin topical solution.

## ADVERSE REACTIONS

**Heartworm-Negative Dogs:** The most common adverse reactions observed during field studies were pruritus, residue, medicinal odor, lethargy, inappetence and hyperactivity.

**Heartworm-Positive Dogs:** The most common adverse reactions observed during field studies were cough, lethargy, vomiting, diarrhea (including hemorrhagic), and inappetence.

## ADVERSE REACTIONS - CATS:

The most common adverse reactions observed during field studies were lethargy, behavioral changes, discomfort, hypersalivation, polydipsia and coughing and gagging. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Vetoquinol USA at 1-800-835-9496. IMOXI™ Topical Solution for Dogs Approved by FDA under ANADA # 200-615. IMOXI™ Topical Solution for Cats Approved by FDA under ANADA # 200-638 Made in U.S.A. © 2019 Vetoquinol USA, Inc. Vetoquinol, the Vetoquinol logo, and IMOXI™ Topical Solution for Dogs and IMOXI™ Topical Solution for Cats are registered trademarks of Vetoquinol USA, Inc. R1 - 05/2020

**Approved for use in dogs and puppies at least 7 weeks of age and weighing at least 3 lbs., and cats and kittens 9 weeks of age or older and weighing at least 2 lbs.**

Disease / Parasite	Dogs and puppies 7 weeks and at least 3 lbs	Cats and kittens 9 weeks and at least 2 lbs
Prevents heartworm disease	X	X
Treats circulating microfilariae	X	
Kills adult fleas + treats flea infestations	X	X
Treats + controls intestinal parasite infections <ul style="list-style-type: none"><li>• Roundworms</li><li>• Hookworms</li><li>• Whipworms</li></ul>	X X X	X X
Treats + controls sarcoptic mange	X	
Treats + controls ear mite infestations		X

**Ask your veterinary team about the benefits of Imoxi™ (imidacloprid + moxidectin) Topical Solution for Dogs and Cats.**

For more information on IMOXI, visit [www.imoxitopical.com](http://www.imoxitopical.com).



**Scan this QR code with your camera to learn more.**



**Scan the QR code to sign up for IMOXI monthly reminders.**